

GAMMA-BAND DESYNCHRONIZATION BY TRANSCUTANEOUS VAGUS NERVE STIMULATION

¹ Department of Otorhinolaryngology—Head and Neck Surgery, HUCH and University of Helsinki
² Department of Biomedical Engineering and Computational Science, Aalto University School of Science
³ School of Medicine, University of Tampere
⁴ BioMag Laboratory, HUS Medical Imaging Center, HUCH and University of Helsinki
⁵ Helsinki Ear Institute

The ability of a treatment method to interfere with tinnitus-related neural activity patterns has been suggested to indicate its potential in relieving tinnitus [1].

Decreased alpha-band (8–12 Hz) and increased gamma-band (> 30 Hz) activity, especially in temporal areas, have been reported in tinnitus patients. Changes in interareal connectivity have also been identified as possible tinnitus markers [2].

Our aim was to investigate the effects of transcutaneous vagus nerve stimulation (tVNS) on evoked and spontaneous auditory cortex activity.

SUBJECTS & METHODS

Table 1 — Characteristic information of the tinnitus patients.

Subject	THI	Age	PTA (L/R) [dB HL]	Tinnitus frequency [Hz]
1	36	30	5 / 2,5	8000
2	58	51	14,4 / 12,5	5000
3	30	47	9,4 / 6,9	1177
4	96	29	5 / 5	7086
5	80	40	13,1 / 5,6	5823
6	50	63	17,5 / 13,1	6000
7	20	29	7,5 / 7,5	4000
Avg.	53	41	10,3 / 7,6	5898

tVNS-induced changes in spectral power and connectivity measures were studied in two scenarios: during presentation of **tinnitus-matched tones** and in **silence**.

Two 5-min measurement blocks were compared: tVNS vs. baseline in both trials. In the active block, tVNS was applied to the left tragus at 25 Hz and approx. 0.25 mA.

Brain activity was registered using a 306-channel whole-head MEG device (Elekta Neuromag). Stimulator artefacts were removed using tSSS [3]. L2 MNE inverse solution was calculated with the dSPM method. Analysis was carried out using MNE-Python [4].

Auditory cortices (ACs) were determined as the Brodmann areas 42 and 43 on each side. Spectral power of ACs and connectivities (squared WPLI) between ACs and other Brodmann areas were calculated.

RESULTS

TINNITUS-MATCHED TONES

The tVNS-induced changes in connectivity are illustrated in **Fig. 1**.

Gamma connectivity decreased in all subjects between left AC and left **BA 32** (*dorsal ACC*, mean decrease 0,021); right AC and right **BA 6** (*premotor cortex*, mean decrease 0,02); and right AC and left **BA 37** (*fusiform gyrus*, mean decrease 0,012).




Figure 1 — tVNS-induced changes in auditory cortex-based connectivity measures during **auditory stimulation**. Blue: decreased connectivity; green: positive correlation with THI; Brown: negative correlation with THI.

Further, the change in **gamma** connectivity between right AC and left **BA 44** (*Broca's area*) correlated negatively with THI scores ($r = -0,964$). Also **beta** connectivity between left AC and right **BA 35** (*perirhinal area*) correlated with THI scores ($r = 0,964$).

THI scores also correlated with the change in left AC **delta** spectral power ($r = 0,964$) and right AC **gamma** spectral power ($r = -0,964$).

SILENCE

The tVNS-induced changes in connectivity are illustrated in **Fig. 2** and given numerically in **Table 2**. THI scores correlated with changes in left AC **alpha** power ($r = -0,893$); **gamma** connectivity between left AC and right **BA 23** (*ventral*

PCC, $r = 0,893$); and in **alpha** connectivity between left AC and left **BA 26** (*cingulate gyrus*, $r = 0,964$).

Table 2 — tVNS-induced changes in auditory cortex-based connectivity measures in silence.

Band	Connection	Mean change
alpha	right AC – right BA 20 (<i>inf. temporal gyrus</i>)	-0,044
	right AC – right BA 21 (<i>middle temporal gyrus</i>)	-0,073
	right AC – right BA 32 (<i>dorsal ACC</i>)	-0,051
	right AC – left BA 29 (<i>retrosplenial cingulate cortex</i>)	-0,033
	right AC – left BA 30 (<i>cingulate cortex</i>)	-0,065
	right AC – left BA 6 (<i>premotor cortex</i>)	+0,021
beta	right AC – right BA 4 (<i>primary motor cortex</i>)	+0,031
	left AC – left BA 17 (<i>primary visual cortex</i>)	+0,020




Figure 2 — tVNS-induced changes in auditory cortex-based connectivity measures in **silence**. Red: increased connectivity; Blue: decreased connectivity; Green: positive correlation with THI.

CONCLUSIONS

tVNS-induced changes were most prominent at gamma band for tone-evoked activity, and at alpha band for spontaneous activity.

tVNS may interfere with dysfunctional auditory-limbic connections found in tinnitus [5].

REFERENCES

- [1] W. Schlee *et al.* 2008. *PLoS ONE* **3**(11):e3720
- [2] N. Weisz & J. Obleser. 2014. *Hear Res* **307**:16–28
- [3] S. Taulu & J. Simola. 2006. *Phys Med Biol* **51**:1759–68
- [4] A. Gramfort *et al.* 2013. *Front Neurosci* **7**:267
- [5] A. M. Leaver *et al.* 2011. *Neuron* **69**:33–43